

Analytically Periodic Solutions to the 3-dimensional Euler-Poisson Equations of Gaseous Stars with Negative Cosmological Constant

YUEN MANWAI*

*Department of Applied Mathematics,
The Hong Kong Polytechnic University,
Hung Hom, Kowloon, Hong Kong*

Revised Date: 11-June-2009

Abstract

By the extension of the 3-dimensional analytical solutions of Goldreich and Weber [6] with adiabatic exponent $\gamma = 4/3$, to the (classical) Euler-Poisson equations without cosmological constant, the self-similar (almost re-collapsing) time-periodic solutions with negative cosmological constant ($\Lambda < 0$) are constructed. The solutions with time-periodicity are novel. On basing these solutions, the time-periodic and almost re-collapsing model is conjectured, for some gaseous stars.

Key Words: Analytically Periodic Solutions, Re-collapsing, Cosmological Constant, Euler-Poisson Equations, Collapsing

1 Introduction

The evolution of a self-gravitating fluid, such as gaseous stars, can be formulated by the isentropic Euler-Poisson systems of the following form:

$$\begin{cases} \rho_t + \nabla \cdot (\rho \vec{u}) = 0, \\ (\rho \vec{u})_t + \nabla \cdot (\rho \vec{u} \otimes \vec{u}) + \nabla P = -\rho \nabla \Phi, \\ \Delta \Phi(t, \vec{x}) = \alpha(N)\rho - \Lambda, \end{cases} \quad (1)$$

where $\alpha(N)$ is a constant related to the unit ball in R^N : $\alpha(2) = 2\pi$ and $\alpha(3) = 4\pi$. And as usual, $\rho = \rho(t, \vec{x})$ and $\vec{u} = \vec{u}(t, \vec{x}) \in \mathbf{R}^N$ are the density, the velocity respectively. $P = P(\rho)$ is the

*E-mail address: nevetsyuen@hotmail.com

pressure function. While Λ is the cosmological constant, for $\Lambda > 0$, the space is open; for $\Lambda < 0$, the space is closed; for $\Lambda = 0$ the space is flat.

In the above systems, the self-gravitational potential field $\Phi = \Phi(t, \vec{x})$ is determined by the density ρ itself, through the Poisson equation (1)₃. For $N = 3$, (1) are the classical (non-relativistic) descriptions of a galaxy, in astrophysics. See [1], [2], [7], for details about the systems.

The γ -law can be applied on the pressure term $P(\rho)$, i.e.

$$P(\rho) = K\rho^\gamma \doteq \frac{\rho^\gamma}{\gamma}, \quad (2)$$

which is a common hypothesis. The constant $\gamma = c_P/c_v \geq 1$, where c_P , c_v are the specific heats per unit mass under constant pressure and constant volume respectively, is the ratio of the specific heats, that is, the adiabatic exponent in (2). In particular, the fluid is called isothermal if $\gamma = 1$. Historically in astrophysics for zero constant ($\Lambda = 0$) background, for the 3-dimensional space, the hydrostatic equilibrium specified by $u = 0$, was studied. According to [2], the ratio between the core density $\rho(0)$ and the mean density $\bar{\rho}$ for $6/5 < \gamma < 2$ is given by

$$\frac{\bar{\rho}}{\rho(0)} = \left(\frac{-3}{z} \dot{y}(z) \right)_{z=z_0}, \quad (3)$$

where y is the solution of the Lane-Emden equation with $n = 1/(\gamma - 1)$ for $\gamma > 1$:

$$\ddot{y}(z) + \frac{2}{z} \dot{y}(z) + y(z)^n = 0, \quad y(0) = \alpha > 0, \quad \dot{y}(0) = 0, \quad n = \frac{1}{\gamma - 1}, \quad (4)$$

and z_0 is the first zero of $y(z_0) = 0$. We can solve the Lane-Emden equation analytically only for

$$y_{anal}(z) \doteq \begin{cases} 1 - \frac{1}{6}z^2, & n = 0; \\ \frac{\sin z}{z}, & n = 1; \\ \frac{1}{\sqrt{1 + z^2/3}}, & n = 5, \end{cases} \quad (5)$$

and for the other values, only numerical values can be obtained. It can be shown that for $n < 5$, the radius of polytropic models is finite; for $n \geq 5$, the radius is infinite.

As we are interested in the solutions in radial symmetry, the Poisson equation (1)₃ is transformed to

$$r^{N-1} \Phi_{rr}(t, x) + (N-1) r^{N-2} \Phi_r = \alpha(N) (\rho - \Lambda) r^{N-1}, \quad (6)$$

$$\Phi_r = \frac{\alpha(N)}{r^{N-1}} \int_0^r (\rho(t, s) - \Lambda) s^{N-1} ds. \quad (7)$$

For the analytical collapsing solutions, the well-known results are constructed by Goldreich and Weber [6] in 1980 for $\gamma = 4/3$ in R^3 . The analytical collapsing solutions without compact support for $\gamma = 1$ in R^2 are constructed by Yuen [11] in 2008.

For simplicity, we take the constant $\Lambda = -\frac{3}{4\pi}$ for $\Lambda < 0$. For the background knowledge for the Euler-Poisson equations with cosmological constant Λ , the interested readers may refer [5], [9].

Our main results are able to obtain the analytically time-periodic solutions for the Euler-Poisson equations with the negative cosmological constant ($\Lambda < 0$) in spherical symmetry:

$$\vec{u} = \frac{\vec{x}}{r} V(t, r), \quad (8)$$

where the diameter $r := \left(\sum_{i=1}^3 x_i^2 \right)^{1/2}$ in R^3 ,

$$\begin{cases} \rho_t + V\rho_r + \rho V_r + \frac{2}{r}\rho V = 0, \\ \rho(V_t + VV_r) + \frac{\partial}{\partial r}(K\rho^{4/3}) = -\frac{4\pi\rho}{r^2} \int_0^r (\rho(t, s) + \frac{3}{4\pi}) s^2 ds, \end{cases} \quad (9)$$

with the extension of the analytical solutions of Goldreich and Weber [6] in the theorem:

Theorem 1 *For the 3-dimensional Euler-Poisson equations in spherical symmetry (9), there exists a family of solutions, for $\gamma = 4/3$:*

$$\begin{cases} \rho(t, r) = \begin{cases} \frac{1}{a(t)^3} y\left(\frac{r}{a(t)}\right)^3, & \text{for } r < a(t)Z_\nu; \\ 0, & \text{for } a(t)Z_\nu \leq r. \end{cases}, \quad V(t, r) = \frac{\dot{a}(t)}{a(t)} r; \\ \ddot{a}(t) = \frac{-\lambda}{a(t)^2} - a(t), \quad a(0) = a_0 > 0, \quad \dot{a}(0) = a_1; \\ \ddot{y}(z) + \frac{2}{z}\dot{y}(z) + \frac{\pi}{K}y(z)^3 = \mu, \quad y(0) = \alpha > 0, \quad \dot{y}(0) = 0, \end{cases} \quad (10)$$

where $K > 0$, $\mu = 3\lambda/(4K)$, and the finite Z_ν is the first zero point of $y(z)$.

In particular,

- (1) $\lambda = 0$ and $a_1 = 0$, the solutions collapse in the finite time $t = \pi/2$;
- (2) $\lambda < 0$, the solutions are non-trivially time-periodic, except the case with $a_0 = \sqrt[3]{\lambda}$ and $a_1 = 0$;
- (3) $\lambda > 0$ and $a_1 \leq 0$, the solutions collapse in a finite time T .

2 Time-Periodic Solutions

In this section, before presenting the proof of Theorem 1, we prepare some lemmas. First, we obtain a general class of solutions for the mass equation in spherical symmetry (9)₁.

Lemma 2 *For the 3-dimensional equation of conservation of mass in spherical symmetry:*

$$\rho_t + V\rho_r + \rho V_r + \frac{2}{r}\rho V = 0, \quad (11)$$

there exist solutions,

$$\rho(t, r) = \frac{f(r/a(t))}{a(t)^3}, \quad V(t, r) = \frac{\dot{a}(t)}{a(t)} r, \quad (12)$$

with the form $f \geq 0 \in C^1$ and $a(t) > 0 \in C^1$.

Proof. We just plug (12) into (11). Then

$$\rho_t + V\rho_r + \rho V_r + \frac{2}{r}\rho V \quad (13)$$

$$= \frac{-3\dot{a}(t)f(r/a(t))}{a(t)^4} - \frac{\dot{a}(t)r\dot{f}(r/a(t))}{a(t)^5} \quad (14)$$

$$+ \frac{\dot{a}(t)r}{a(t)} \frac{\dot{f}(r/a(t))}{a(t)^4} + \frac{f(r/a(t))}{a(t)^3} \frac{\dot{a}(t)}{a(t)} + \frac{2}{r} \frac{f(r/a(t))}{a(t)^3} \frac{\dot{a}(t)}{a(t)} r \quad (15)$$

$$= 0. \quad (16)$$

The proof is completed. ■

The following core lemma is needed to show the cyclic phenomena of the solutions (10).

Lemma 3 *For the ordinary differential equation,*

$$\ddot{a}(t) - \frac{\lambda}{a(t)^2} + a(t) = 0, \quad a(0) = a_0 > 0, \quad \dot{a}(0) = a_1, \quad (17)$$

with $\lambda > 0$, there exist non-trivially periodic solutions except the case with $a_0 = \sqrt[3]{\lambda}$ and $a_1 = 0$.

Proof. To the equation (17), multiply $\dot{a}(t)$ and then integrate it:

$$\frac{\dot{a}(t)^2}{2} + \frac{\lambda}{a(t)} + \frac{a(t)^2}{2} = \theta, \quad (18)$$

with the constant $\theta = \frac{a_1^2}{2} + \frac{\lambda}{a_0} + \frac{a_0^2}{2} > 0$.

We define the kinetic energy:

$$F_{kin} := \frac{\dot{a}(t)^2}{2}, \quad (19)$$

and the potential energy:

$$F_{pot} = \frac{\lambda}{a(t)} + \frac{a(t)^2}{2}. \quad (20)$$

The total energy is conserved:

$$\frac{d}{dt}(F_{kin} + F_{pot}) = 0. \quad (21)$$

Observe the potential energy has only one global minimum at $a_{\min} = \sqrt[3]{\lambda}$, for $a(t) \in (0, +\infty)$.

Therefore, by the classical energy method for conservative systems (in Section 4.3 of [8]), the solutions have the closed trajectory. We calculate the time for traveling the closed orbit:

$$T = \int_0^t dt = 2 \int_{a_{\max}}^{a_{\min}} \frac{-da(t)}{\sqrt{2[\theta - (\frac{\lambda}{a(t)} + \frac{a(t)^2}{2})]}} = 2 \int_{a_{\min}}^{a_{\max}} \frac{da(t)}{\sqrt{2[\theta - (\frac{\lambda}{a(t)} + \frac{a(t)^2}{2})]}}, \quad (22)$$

where $a_{\min} = \inf_{t \geq 0} (a(t))$ and $a_{\max} = \sup_{t \geq 0} (a(t))$.

Let $G(t) := \theta - (\frac{\lambda}{a(t)} + \frac{a(t)^2}{2})$, $G_0 := \theta - (\frac{\lambda}{a_{\min} + \epsilon} + \frac{(a_{\min} + \epsilon)^2}{2}) > 0$ and $G_1 := \theta - (\frac{\lambda}{a_{\min} + \epsilon} + \frac{(a_{\min} + \epsilon)^2}{2}) > 0$.

Except the case with $a_0 = \sqrt[3]{\lambda}$ and $a_1 = 0$, the equation (22) becomes

$$T = \int_{a_{\min}}^{a_{\min}+\epsilon} \frac{2da(t)}{\sqrt{2[\theta - (\frac{\lambda}{a(t)} + \frac{a(t)^2}{2})]}} + \int_{a_{\min}+\epsilon}^{a_{\max}-\epsilon} \frac{2da(t)}{\sqrt{2[\theta - (\frac{\lambda}{a(t)} + \frac{a(t)^2}{2})]}} \quad (23)$$

$$+ \int_{a_{\max}-\epsilon}^{a_{\max}} \frac{2da(t)}{\sqrt{2[\theta - (\frac{\lambda}{a(t)} + \frac{a(t)^2}{2})]}} \quad (24)$$

$$\leq \sup_{a_{\min} \leq a(t) \leq a_{\min}+\epsilon} \left| \frac{1}{\frac{\lambda}{a(t)^2} - a(t)} \right| \int_0^{G_0} \frac{\sqrt{2}dG(t)}{\sqrt{G(t)}} + \int_{a_{\min}+\epsilon}^{a_{\max}-\epsilon} \frac{2da(t)}{\sqrt{2[\theta - (\frac{\lambda}{a(t)} + \frac{a(t)^2}{2})]}} \quad (25)$$

$$+ \sup_{a_{\max}-\epsilon \leq a(t) \leq a_{\max}} \left| \frac{1}{\frac{\lambda}{a(t)^2} - a(t)} \right| \int_0^{G_1} \frac{\sqrt{2}dG(t)}{\sqrt{G(t)}} \quad (26)$$

$$= \sup_{a_{\min} \leq a \leq a_{\min}+\epsilon} \left| \frac{1}{\frac{\lambda}{a(t)^2} - a(t)} \right| \frac{\sqrt{G_0}}{\sqrt{2}} + \int_{a_{\min}+\epsilon}^{a_{\max}-\epsilon} \frac{2da(t)}{\sqrt{2[\theta - (\frac{\lambda}{a(t)} + \frac{a(t)^2}{2})]}} \quad (27)$$

$$+ \sup_{a_{\max}-\epsilon \leq a(t) \leq a_{\max}} \left| \frac{1}{\frac{\lambda}{a(t)^2} - a(t)} \right| \frac{\sqrt{G_1}}{\sqrt{2}} \quad (28)$$

$$< \infty. \quad (29)$$

Therefore, the solutions to the differential equation (17) are time-periodic.

The proof is completed ■

In the solutions (10), due to the properties of the modified Emden equation (10)₂, the collapsing phenomena are observed. The following lemma can be proved immediately.

Lemma 4 *For the ordinary differential equation,*

$$\ddot{a}(t) + \frac{\lambda}{a(t)^2} = -a(t), \quad 0 < a(0) = a_0, \quad \dot{a}(0) = a_1 \leq 0, \quad (30)$$

with $\lambda > 0$,

there exists a finite time T such that $\lim_{t \rightarrow T^-} a(t) = 0$.

Proof. If the claim is not true, for all $t \geq 0$, we have $a(t) > 0$.

But the ordinary differential equation (30) becomes

$$\ddot{a}(t) = \frac{-1}{a(t)^2} - a(t) < 0. \quad (31)$$

This shows $\dot{a}(t)$ is decreasing function: $\dot{a}(t) < \dot{a}(t_1)$ for all $t > t_1 > 0$,

$$\dot{a}(t) \leq \dot{a}(t_1) < 0. \quad (32)$$

Thus, the solution is bounded by

$$a(t) = \int_{t_1}^t \dot{a}(s)ds + a(t_1) \leq \dot{a}(t_1)t + a_0. \quad (33)$$

After a sufficient large time, there exists a finite time T such that $\lim_{t \rightarrow T^-} a(t) = 0$. A contraction is met.

The proof is completed. ■

Here we are ready to give proof of Theorem 1.

Proof of Theorem 1. From Lemma 2, we can easily check that (10) satisfy (9)₁. We plug the solutions (10), into the momentum equation (9)₂:

$$\rho(V_t + VV_r) + K \frac{\partial}{\partial r} \rho^{4/3} + \frac{4\pi\rho}{r^2} \int_0^r (\rho(t, s) - \Lambda) s^2 ds \quad (34)$$

$$= \rho \frac{\ddot{a}(t)}{a(t)} r + 4K \left(\frac{y(\frac{r}{a(t)})^3}{a(t)^3} \right)^{1/3} \frac{y(\frac{r}{a(t)})^2 \dot{y}(\frac{r}{a(t)})}{a(t)^4} + \frac{4\pi\rho}{r^2} \int_0^r (\rho(t, s) - \Lambda) s^2 ds \quad (35)$$

$$= \rho \left[\frac{-\frac{\lambda}{a(t)^2} + \frac{4\pi\Lambda}{3} a(t)}{a(t)} r \right] + 4K \frac{y(\frac{r}{a(t)})^3}{a(t)^3} \frac{\dot{y}(\frac{r}{a(t)})}{a(t)^2} + \frac{4\pi\rho}{r^2 a(t)^2} \int_0^r y^3 \left(\frac{s}{a(t)} \right) s^2 ds - \frac{4\pi\Lambda\rho}{r^2} \int_0^r s^2 ds \quad (36)$$

$$= \rho \frac{-\lambda r}{a(t)^3} + \frac{4\pi\Lambda}{3} \rho r + 4K \rho \frac{\dot{y}(\frac{r}{a(t)})}{a(t)^2} + \frac{4\pi\rho}{r^2 a(t)^3} \int_0^r y \left(\frac{s}{a(t)} \right)^3 s^2 ds - \frac{4\pi\Lambda}{3} \rho r \quad (37)$$

$$= \frac{\rho}{a(t)^2} \left[-\frac{\lambda}{a(t)} r + 4K \dot{y} \left(\frac{r}{a(t)} \right) + \frac{4\pi}{r^2 a(t)} \int_0^r y \left(\frac{s}{a(t)} \right)^3 s^2 ds \right] \quad (38)$$

$$= \frac{\rho}{a(t)^2} \left[-\frac{\lambda}{a(t)} r + 4K \dot{y} \left(\frac{r}{a(t)} \right) + \frac{\alpha(N)}{(\frac{r}{a(t)})^2} \int_0^{r/a(t)} y(s)^3 s^2 ds \right] \quad (39)$$

$$= \frac{\rho}{a(t)^2} Q \left(\frac{r}{a(t)} \right). \quad (40)$$

Here, we use the property of $a(t)$:

$$\ddot{a}(t) = \frac{-\lambda}{a(t)^2} + \frac{4\pi\Lambda}{3} a(t) = \frac{-\lambda}{a(t)^2} - a(t), \quad (41)$$

and denote

$$Q\left(\frac{r}{a(t)}\right) := Q(x) = -\lambda x + 4K \dot{y}(x) + \frac{4\pi}{x^2} \int_0^x y(s)^3 s^2 ds. \quad (42)$$

Differentiate $Q(x)$ with respect to x ,

$$\dot{Q}(x) = -\lambda + 4K \ddot{y}(x) + 4\pi y(x)^3 - \frac{2 \cdot 4\pi}{x^3} \int_0^x y(s)^3 s^2 ds \quad (43)$$

$$= -\frac{2}{x} \left[\lambda x + 4K \dot{y}(x) - K \mu x + \frac{4\pi}{x^2} \int_0^x y(s)^3 s^2 ds \right] \quad (44)$$

$$= -\frac{2}{x} Q(x). \quad (45)$$

With $\lim_{x \rightarrow 0^+} Q(x) = Q(0) = 0$, this implies that $Q(x) = 0$.

The above result is due to the fact that we choose the Lane-Emden equation:

$$\begin{cases} \ddot{y}(z) + \frac{2}{z} \dot{y}(z) + \frac{\pi}{K} y(z)^3 = \mu, \mu = \frac{3\lambda}{4K}, \\ y(0) = \alpha > 0, \dot{y}(0) = 0. \end{cases} \quad (46)$$

We have shown that there exists the family of the solutions with compact support, in Theorem 1, with the well-known results about the Lane-Emden equation (46) [2] [7].

On the other hand, for $\Lambda = -\frac{3}{4}\pi < 0$, it is clear for $\lambda = 0$ in the theorem is true. For the linear ordinary differential equation:

$$\ddot{a}(t) + a(t) = 0, \quad a(0) = a_0 > 0, \quad \dot{a}(0) = 0, \quad (47)$$

the solution is

$$a(t) = a_0 \cos t. \quad (48)$$

The function $a(\frac{\pi}{2})$ achieves zero and the solutions collapse in the finite time $\pi/2$.

By using Lemma 3 about $a(t)$, we have, for $\lambda < 0$, the periodic solutions are non-trivial except the case with $a_0 = \sqrt[3]{\lambda}$ and $a_1 = 0$.

By using Lemma 4, we have, for $\lambda > 0$ and $a_1 \leq 0$ the solutions collapse in a finite time T .

This completes the proof. ■

Remark 5 *If we consider the system with frictional damping,*

$$\begin{cases} \rho_t + \nabla \bullet (\rho \vec{u}) = 0, \\ (\rho \vec{u})_t + \nabla \bullet (\rho \vec{u} \otimes \vec{u}) + \beta \rho \vec{u} + \nabla P = -\rho \nabla \Phi, \\ \Delta \Phi(t, \vec{x}) = \alpha(N)\rho - \Lambda, \end{cases} \quad (49)$$

with the constant $\beta > 0$,

the corresponding ordinary differential equation for the solutions are:

$$\ddot{a}(t) + \beta \dot{a}(t) = \frac{-\lambda}{a(t)^2} - a(t), \quad a(0) = a_0 > 0, \quad \dot{a}(0) = a_1, \quad (50)$$

[10].

For $\lambda < 0$, the function $a(t)$ oscillates around the minimum $a_{\min} = \sqrt[3]{\lambda}$ of the potential energy except the case with $a_0 = \sqrt[3]{\lambda}$ and $a_1 = 0$. And it is asymptotically stable.

Remark 6 *In general, for the Euler-Poisson equations in R^N , it is clear to have the corresponding solutions:*

for $N = 2$ and $\gamma = 1$,

$$\begin{cases} \rho(t, r) = \frac{1}{a^2(t)} e^{y(r/a(t))}, \quad V(t, r) = \frac{\dot{a}(t)}{a(t)} r; \\ \ddot{a}(t) = \frac{-\lambda}{a(t)} - a(t), \quad a(0) = a_0 > 0, \quad \dot{a}(0) = a_1; \\ \ddot{y}(z) + \frac{1}{z} \dot{y}(z) + \frac{2\pi}{K} e^{y(z)} = \mu, \quad y(0) = \alpha, \quad \dot{y}(0) = 0, \end{cases} \quad (51)$$

where $\mu = 2\lambda/K$, [11];

for $N \geq 3$ and $\gamma = (2N - 2)/N$,

$$\left\{ \begin{array}{l} \rho(t, r) = \begin{cases} \frac{1}{a^N(t)} y\left(\frac{r}{a(t)}\right)^{N/(N-2)}, & \text{for } r < a(t)Z_\mu; \\ 0, & \text{for } a(t)Z_\mu \leq r. \end{cases}, \quad V(t, r) = \frac{\dot{a}(t)}{a(t)}r, \\ \ddot{a}(t) = \frac{-\lambda}{a^{N-1}(t)} - a(t), \quad a(0) = a_0 > 0, \quad \dot{a}(0) = a_1, \\ \ddot{y}(z) + \frac{N-1}{z}\dot{y}(z) + \frac{\alpha(N)}{(2N-2)K}y(z)^{N/(N-2)} = \mu, \quad y(0) = \alpha > 0, \quad \dot{y}(0) = 0, \end{array} \right. \quad (52)$$

where $\mu = [N(N-2)\lambda]/(2N-2)K$ and the finite Z_μ is the first zero of $y(z)$ [4].

With $\lambda < 0$, the solutions 51 and 52 are time-periodic for the negative cosmological constant.

Remark 7 Under some initial conditions for $\Lambda > 0$, the collapsing solutions are obtained. The case is similar to the system without cosmological constant.

3 Discussion

By the separation method, the solutions are constructed in its close relatives (the Euler-Poisson equations with frictional damping, Navier-Stokes, Euler equations and the ones with frictional damping term) [10] and [12]. However, we emphasis on that it is novel to have the time-periodic pattern in the solution structure with $\Lambda < 0$.

On the other hand, in the statement (2) of the theorem, with some initial conditions, for example, $0 < a(0) = \epsilon < 1$ and $\dot{a}(0) = 0$, the density becomes

$$\rho(0, 0) = \frac{\alpha^3}{\epsilon^3} \gg M_0, \quad (53)$$

where M_0 is an arbitrary constant.

It may provide the possible evolutionary model with $\Lambda < 0$ constant in mathematical theory, that the universe expands and then almost re-collapses (The density at the origin can be greater than any given constant $\rho(T, 0) \gg M_0$), in the finite time periodically. The solutions may be like a black hole (in the sense, $\rho(T, 0) = +\infty$ with a finite time T). The black hole seems to be the end of the evolution of the periodic solutions. However actually, there exists no authentic black hole in our particular time-periodic solutions.

As the time-periodic effect is due to the negative cosmological constant in our model, more solutions with this pattern are expected for other γ values. Then further works may be done by the computing simulations about the stability of the numerical solutions. If some gaseous stars (a galaxy) obey the γ -law ($\gamma = 4/3$), it may provide an alternative explanation about that, the time-periodic solutions coincide with the expansion segment (the red-shift effect) in a short time. We notice that it is the significant difference between the traditional re-collapsing model with negative cosmological constant $\Lambda < 0$ [3].

References

- [1] J. Binney and S. Tremaine, *Galactic Dynamics*, Princeton Univ. Press, 1994.
- [2] S. Chandrasekhar, *An Introduction to the Study of Stellar Structure*, Univ. of Chicago Press, 1939.
- [3] M.P. Dąbrowski, *Future State of the Universe*, (2006) Annalen der Physik (Leipzig), 15 (4-5), pp. 352-363.
- [4] Y.B. Deng, J.L. Xiang and T. Yang, *Blowup Phenomena of Solutions to Euler-Poisson Equations*, J. Math. Anal. Appl. **286** (1)(2003), 295-306.
- [5] H. H. Fliche and R. Triay, *Euler-Poisson-Newton Approach in Cosmology*, Cosmology and Gravitation, 346–360, AIP Conf. Proc., 910, Amer. Inst. Phys., Melville, NY, 2007.
- [6] P. Goldreich and S. Weber, *Homologously Collapsing Stellar Cores*, Astrophys. J. **238**, 991 (1980).
- [7] R. Kippenhahn and A. Weigert, *Stellar Structure and Evolution*, Springer-Verlag, 1990.
- [8] W.D. Lakin and D. A. Sanchez, *Topics in Ordinary Differential Equations*, Dover Pub. Inc., New York, 1982.
- [9] P.J.E. Peebles and B. Ratra, *The cosmological constant and dark energy*, Reviews of Modern Physics, Volume 75, Issue 2, April 2003, Pages 559-606.
- [10] M.W. Yuen, *Blowup Solutions for a Class of Fluid Dynamical Equations in R^N* , J. Math. Anal. Appl. **329** (2)(2007), 1064-1079.
- [11] M.W. Yuen, *Analytical Blowup Solutions to the 2-dimensional Isothermal Euler-Poisson Equations of Gaseous Stars*, J. Math. Anal. Appl. **341** (1)(2008), 445-456.
- [12] M.W. Yuen, *Analytical Solutions to the Navier-Stokes Equations*, J. Math. Phys., **49** (2008) No. 11, 113102, 10pp.